

Vaccination Schedule 2021

More than
10 YEARS
of Excellence

Disease	Vaccine	Birth	6 weeks	10 weeks	14 weeks	6 months	9 months	12 months	15 months	18 months	6 years	9 years	12 years
TB	BCG	●											
Polio	bOPV	●	●										
Diphtheria, Tetanus, Polio, Pertussis Haemophilus Influenzae, Hepatitis B	Hexaximi		●	●	●					●			
	Infanrix Hexa		8 weeks ●	12 weeks ●	16 weeks ●					●			
Pneumococcal	Prevenar 13		●		●		●						
	Synflorix		●	●	●		●	●	Nine and a half months				
Rotavirus	Rotarix		●		●								
	RotaTeq		●	●	●								
Measles	Measbio					●		●	12 months Measbio not required if giving MMR				
Measles Mumps Rubella	Priorix							●	MMR at 12 months instead of Measbio		●	Give MMR and Varilrix or Onvara separately at 6 years if not using Priorix Tetra	
	MSD OMZYTA							●			●		
Chickenpox	Varilrix								●		●		
	ONVARA								●		●		
Measles Mumps Rubella + Chickenpox	Priorix Tetra	Priorix Tetra can be used at 12 months instead of Measbio, MMR and Varicella. Increased risk of febrile convulsion after the first dose of Priorix Tetra						●			●		
Meningococcal Conjugate	Menactra						●		●	For individuals 2 - 55 years administer 1 dose			
Hepatitis A	AVAXIM ⁸⁰							●	Hep A with MMR not with Measbio		●		
	Havrix Junior							●			●		
Tetanus, Diphtheria Pertussis, Polio	TETRA XIM								Tetraxim at 6 years or Boostrix Tetra at 6 years Boostrix Tetra can be given at 12 years if not given at 6 years. Adacel Quadra at 12 years		●		
	Boostrix Tetra										●		
	ADACEL QUADRA												●
Tetanus, Diphtheria	Td										●		●
Human Papilloma Virus	Cervarix	Cervarix from 9 -14 years or Gardasil from 9 -13 years as 2 doses 6 months apart State programme: Cervarix to Grade 4 girls in public schools 9 -14 years. 2 doses 6 months apart										●	
	GARDASIL											●	
Influenza	Influvac	From 6 months, important for children in creche, chronic illness and respiratory problems. Start before the INFLUENZA season in MAY.											
	Vaxigrip Tetra												
Respiratory Syncytial Virus	SYNAGIS [®] PALIVIZUMAB	RSV prophylaxis in high-risk infants-prevention of serious LRTI caused by RSV. Start January and end in May.											
Pneumococcal	PNEUMOVAX ²³ (pneumococcal vaccine, polyvalent, MSD)	Must have at least one dose of Conjugate PCV before Pneumovax 23. Pneumovax only in children older than 2 years with Immune Compromise or high risk of Pneumococcal infection. Two doses with 2nd dose 3 - 5 years after first.											

● **STATE EPI VACCINES** These vaccines are available free from Government supplied clinics

● **RECOMMENDED OPTIONAL VACCINES** Some of the vaccines in this schedule are only available from private clinics